

## WO9623000

Publication Title:

Nucleic acids encoding ligands for HEK4 receptors

Abstract:

Polypeptides which bind to one or more EPH-like receptors, particularly the HEK4 receptor, are described. The polypeptides are designated HEK4 binding proteins. Nucleic acids encoding HEK4 binding proteins, and expression vectors, host cells and processes for the production of the polypeptides are also described. The polypeptides are useful for modulating the growth and/or differentiation of a variety of tissues, including those from liver, kidney, lung, skin, digestive tract and nervous system and may be used to regenerate damaged or depleted tissue and to treat cancer or nervous system disorders.

-----  
Data supplied from the esp@cenet database - <http://ep.espacenet.com>

*This Patent PDF Generated by Patent Fetcher(TM), a service of Stroke of Color, Inc.*



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C07K 14/47, C12N 15/12, C07K 16/18, C12N 15/70, G01N 33/68, C12Q 1/68, A61K 38/17, C07K 19/00</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 96/23000</b> <b>(43) International Publication Date:</b> 1 August 1996 (01.08.96)
<b>(21) International Application Number:</b> PCT/US96/01079 <b>(22) International Filing Date:</b> 16 January 1996 (16.01.96)  <b>(30) Priority Data:</b> 08/379,802 27 January 1995 (27.01.95) US  <b>(71) Applicant:</b> AMGEN INC. [US/US]; Amgen Center, 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).  <b>(72) Inventors:</b> BARTLEY, Timothy, D.; 2431 McCrea Road, Thousand Oaks, CA 91362 (US). FOX, Gary, M.; 35 West Kelly Road, Newbury Park, CA 91320 (US).  <b>(74) Agents:</b> ODRE, Steven, M. et al.; Amgen Inc., Amgen Center, 1840 Dehavilland Drive, Thousand Oaks, CA 91320-1789 (US).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AZ, BY, KG, KZ, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
<b>(54) Title:</b> LIGANDS FOR EPH-LIKE RECEPTORS  <b>(57) Abstract</b>  Polypeptides which bind to one or more EPH-like receptors, particularly the HEK4 receptor, are described. The polypeptides are designated HEK4 binding proteins. Nucleic acids encoding HEK4 binding proteins, and expression vectors, host cells and processes for the production of the polypeptides are also described. The polypeptides are useful for modulating the growth and/or differentiation of a variety of tissues, including those from liver, kidney, lung, skin, digestive tract and nervous system and may be used to regenerate damaged or depleted tissue and to treat cancer or nervous system disorders.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

## LIGANDS FOR EPH-LIKE RECEPTORS

The invention relates to polypeptides which bind to one or more EPH-like receptors. More particularly, the invention relates to polypeptides which bind to the HEK4 receptor, to nucleic acids encoding same and to expression vectors and host cells for the production of the polypeptides.

10 Background of the Invention

The response of cells to their environment is often mediated by soluble protein growth and differentiation factors. These factors exert their effects by binding to and activating transmembrane receptors. This interaction is the initial event in a cascade which culminates in a biological response by the cell. An important class of transmembrane receptors is the receptor protein tyrosine kinases (receptor PTKs, reviewed in van der Geer et al. Ann. Rev. Cell. Biol. 10, 251-337 (1994). PTKs consist of an extracellular domain which interacts specifically with the receptor's cognate ligand, a membrane spanning domain, and an intracellular domain which harbors the tyrosine kinase activity. Receptor PTKs are activated by ligand-mediated dimerization followed by autophosphorylation of tyrosine residues in the cytoplasmic domain. The receptor PTK can then in turn phosphorylate substrate molecules in the signal transduction pathway, leading to a cellular response.

The family of receptor PTKs can be divided into a number of sub-families based on the general structure of the extracellular domain and on amino acid sequence relationships within the catalytic domain. Currently, the largest known sub-family of

- 2 -

receptor protein tyrosine kinases is the EPH-like receptors, consisting of at least 13 members. Members of this sub-family include the following: EPH (Hirai et al., Science 238, 1717-1725 (1987)), ECK (Lindberg et al., Mol. Cell. Biol. 10, 6316-6324 (1990)), Cek4, 5 Cek5, Cek6, Cek7, Cek8, Cek9, Cek10 (Pasquale, Cell Regulation 2, 523-534 (1991); Sajjadi et al., The New Biologist 3, 769-778 (1991); Sajjadi and Pasquale, Oncogene 8, 1807-1813 (1993)), Eek, Erk (Chan and 10 Watt, Oncogene 6, 1057-1061 (1991)), Ehk1, Ehk2 (Maisonpierre et al., Oncogene 8, 3277-3288 (1993)), HEK (PCT Application No. W093/00425; Wicks et al., PNAS 89, 1611-1615 (1992)), HEK2 (Bohme et al., Oncogene 8, 2857-2862 (1993)), HEK5, HEK7, HEK8, HEK11 15 (U.S. Serial No. 08/229,509) and HTK (Bennett et al. J. Biol. Chem. 269, 14211-14218 (1994)).

Until recently, no ligands for any member of the EPH sub-family had been identified. A ligand for the Eck receptor was described in PCT Application 20 No. WO 94/11020 and Bartley et al. (Nature 368, 558-560 (1994)) and identified earlier as B61, a polypeptide encoded by a cDNA of unknown function (Holzman et al., Mol. Cell Biol. 10, 5830-5838 (1990)). Ligands for Elk and Ehk1 receptors have also 25 been reported (PCT Application No. W094/11384; Davis et al., Science 266, 816-819 (1994)). Most recently, a polypeptide (ELF-1) identified from a mouse embryo midbrain and hindbrain cDNA library has been reported to be a ligand for Mek4 and Sek (Cheng and Flanagan, 30 Cell 79, 157-168 (1994)).

Most attempts to purify soluble factors from complex biological fluids have depended on cell-based bioassays of the response to stimulation by the factor. These include increased cell growth or 35 survival, increased DNA synthesis, a chemotactic response, or some other downstream consequence of

- 3 -

receptor activation. Receptor autophosphorylation has also been used as an assay to detect stimulation of the cell. We have recently described a method for the isolation of ligands based on direct detection of  
5 receptor/ligand binding and the use of receptor affinity chromatography for purification (Bartley et al., supra). Here we report the application of this method to purify, sequence, and molecularly clone one of a family of ligands corresponding to the EPH sub-  
10 family of receptor tyrosine kinases.

Although the EPH sub-family is the largest known sub-family of receptor PTKs, few ligands have been identified which bind to and activate an EPH sub-family receptor. It is therefore an objective to  
15 identify additional ligands for EPH sub-family receptor PTKs. These ligands will be useful for modulating responses of EPH sub-family receptor bearing cells.

## 20 Summary of the Invention

The present invention relates to polypeptides capable of binding to one or more EPH-like receptor PTKs. More particularly, the invention  
25 provides polypeptides which bind to the HEK4 receptor, but may also bind to other members of the sub-family of EPH-like receptor PTKs. These polypeptides are referred to as HEK4 binding proteins (HEK4 BPs). In one embodiment, the polypeptide binds to and activates  
30 HEK4 and ECK receptors. Also encompassed by the invention are nucleic acids encoding HEK4 BPs and nucleic acids which hybridize to HEK4 BP nucleic acids and encode polypeptides having at least one of the biological properties of a HEK4 BP. Biologically  
35 active HEK4 BP fragments and analogs and nucleic acids

- 4 -

encoding same as well as fusion proteins comprising HEK4 BP are also encompassed by the invention.

Expression vectors and host cells for the production of biologically active HEK4 BP and processes for the production of HEK4 BP using the expression vectors and host cells are also within the scope of the invention. Antibodies specifically binding HEK4 BP are also provided for.

Polypeptides of the invention are useful for modulating (i.e., increasing or decreasing) the growth and/or differentiation of EPH sub-family receptor-bearing cells, particularly cells expressing HEK4 or ECK receptors. Based on levels of expression of HEK4, ECK, and HEK4 BP in a variety of tissues, it is expected that HEK4 BP will be useful for modulating the growth and/or differentiation, for example, liver, kidney, lung, skin or neural tissues. Administration of HEK4 BP to mammals is useful in the treatment of nervous system disorders and in the regeneration of damaged or depleted tissues. HEK4 BP antagonists are also useful for the treatment of cancers.

#### Description of the Figures

Figure 1. BIAcore screening of conditioned media on HEK4-X surface. Concentrated samples of cell-conditioned media were screened on a HEK4-X surface as described in Example 2. The number of conditioned media samples giving a signal within each range of resonance units (RU) are shown in the histogram. Samples which bound more than 200 RU are summarized in Table 1.

Figure 2. Purification of HEK4 Binding Protein from A498 conditioned media. C4 Reverse Phase

- 5 -

HPLC column profile of HEK4 BP (a); SDS-PAGE analysis of pools of indicated peaks observed on the C4 column.

Figure 3. Sequence of HEK-4 binding protein cDNA. The nucleic acid sequence of the human HEK-4 binding protein cDNA clone containing the entire coding sequence is shown along with the predicted amino acid sequence. The cDNA clone predicts a protein of between 213 and 228 amino acids, depending on which of three potential start codons is utilized. The sequence is numbered so that the predicted mature N-terminal amino acid is residue 1, with the putative signal peptide (underlined) extending from residues -19 to -1.

Figure 4. Purification of recombinant HEK-4 binding protein. C4 Reverse Phase HPLC column profile of recombinant HEK4 BP (a); SDS-PAGE analysis of C4 fractions in the vicinity of the A<sub>214</sub> peak. Fractions are identified by elution times from the C4 column.

Figure 5. Expression of HEK-4 binding protein in human tissues. The expression of HEK-4 binding protein mRNA in human tissues was examined by Northern blot analysis as described in Example 6. A blot containing 2 µg of polyA<sup>+</sup> mRNA isolated from each of several tissues was purchased from Clontech (Palo Alto, CA) and hybridized with a <sup>32</sup>P-labeled HEK-4 binding protein cDNA probe.

Figure 6. Stimulation of tyrosine phosphorylation of EPH-like receptors by membrane-bound HEK4 BP. CHO cells that express recombinant HEK4 receptor and endogenous ECK were treated with cells that were transfected with an



- 6 -

expression vector that contained the HEK4 BP cDNA or vector without cDNA. After lysis, HEK4 receptor (a) or ECK receptor (b) were immunoprecipitated. The immunoprecipitates were fractionated by PAGE, electrophoretically transferred, and probed with antiphosphotyrosine antibodies.

Figure 7. Stimulation of tyrosine phosphorylation by soluble HEK4 BP. Cells were treated with conditioned media (CM) or recombinant HEK4 BP, with (+) or without (-) antibody clustering, and assayed for HEK4 receptor activation. a) Twelve-fold concentrated media was compared to 2µg/ml HEK4 BP. (b) A dose response comparing clustered and unclustered HEK4 BP.

Figure 8. Relative affinity of HEK4 BP for HEK 4, ECK and HEK8 receptors. A competition assay for measuring binding of HEK4 BP to immobilized HEK4 receptor in the presence of increasing concentrations of soluble HEK4, ECK and HEK8 receptors was performed as described in Example 8. The line identified as "IC 50" is drawn at an RU value corresponding to a HEK4 BP concentration which is 50% of the control (no competitor) RU value.

#### Detailed Description of the Invention

The present invention relates to polypeptides capable of binding to one or more EPH-like receptor PTKs and, more particularly, are capable of binding to a human homolog of an EPH-like receptor PTK. To date, eight human homologs of EPH-like receptors have been identified: EPH, ECK, HTK, HEK2, HEK4, HEK5, HEK7, HEK8 and HEK11. Characteristics of several HEK receptors are disclosed

- 7 -

in co-pending and commonly owned U.S. Serial No. 08/229,509 herein incorporated by reference. The polypeptides of the present invention preferentially bind HEK4 receptor and are referred to herein as HEK4  
5 binding proteins (HEK4 BP). The HEK4 receptor is a glycosylated 135 kDa protein tyrosine kinase previously identified as HEK receptor by Wilks et al. supra, and is the human homolog of the Cek 4 and Mek 4 receptors identified in chicken and mouse,  
10 respectively. Polypeptides capable of binding HEK4 receptor may also activate the receptor by inducing receptor autophosphorylation, an event which initiates transmission of a signal from the cell surface to the nucleus. Activation of HEK4 receptors leads to  
15 modulation of growth and/or differentiation of HEK4 receptor-bearing cells. HEK4 BP also binds to and activates other EPH-like receptors as described below.

A HEK4 binding protein has been identified  
20 and isolated from A498 cell line conditioned medium by procedures generally described in U.S. Serial No. 08/145,616, relevant portions of which are herein incorporated by reference, and described in more detail in Example 2. Briefly, a gene encoding the  
25 extracellular domain of the HEK4 was constructed and expressed as described in Example 1. The purified HEK4 extracellular domain was immobilized on a BIAcore sensor chip and concentrated conditioned media from 102 different cell lines were screened for binding to  
30 HEK4 receptor extracellular domains by surface plasmon resonance. This procedure identified conditioned medium from several cell lines shown in Table 1 having one or more factors which interacted with HEK4. The A498 cell line was chosen as a source for HEK4 ligand  
35 and a protein binding to HEK4 was purified as described in Example 2. Purified and isolated HEK4

- 8 -

binding protein from A498 cell-conditioned medium has three major forms of molecular weights 21, 25 and 27kD on nonreducing SDS-polyacrylamide gels. These forms represent glycosylation and C- terminal processing variants. HEK4 BP has the amino acid sequences as shown in Table 2 for the peptides generated by cyanogen bromide or trypsin cleavage.

cdNA clones of HEK4 binding protein were obtained from a human placenta cdNA library as described in Example 3. The sequence of human HEK4 binding protein cdNA is shown in Figure 3. Based upon cdNA sequencing and carboxy-terminal peptide mapping of the A498 cell-derived protein, a major secreted form of HEK4 binding protein had an amino terminal serine residue as shown in Figure 3 and a carboxy terminal proline residue at position 179. An alternate secreted form having a carboxy terminal alanine residue at position 177 was also detected. Alternative forms of HEK4 BP, including membrane-bound forms, may also be synthesized.

Recombinant HEK4 binding protein was expressed in CHO cells transfected with cdNA encoding HEK4 BP as shown in Figure 3. Soluble HEK4 BP was purified as described in Example 4 and shown to bind HEK4 receptor by BIAcore analysis. Purified soluble HEK4 BP activated HEK4 receptor and this activation was enhanced by antibody clustering of the ligand (Figure 7). Activation of HEK4 receptor was also observed upon contact of CHO cells expressing HEK4 BP with CHO cells expressing HEK4 receptor (Figure 6). Therefore, recombinant HEK4 binding protein binds and activates EPH-like receptor PTKs.

The invention provides for a purified and isolated polypeptide, termed HEK4 binding protein,

- 9 -

capable of binding at least one EPH-like receptor. In one embodiment, the polypeptide is capable of binding the HEK4 receptor. HEK4 binding protein is mammalian and is preferably human. Purified HEK4 binding protein is substantially free of other human proteins and has a molecular weight of about 21 to 27kD on nonreducing SDS-PAGE. HEK4 BP has at least about 70% homology to the amino acid sequence as shown in Figure 3 (SEQ ID NO: 1) and is capable of binding at least one EPH-like receptor. Preferably, HEK4 BP has the amino acid sequence as shown in Figure 3 (SEQ ID NO: 1). Binding of an EPH-like receptor by HEK4 BP may or may not result in receptor activation. EPH-like receptor binding and activation may be effected by either soluble or membrane-bound form of HEK4 BP, or by both forms. It is further understood that receptor binding and activation by HEK4 BP is not restricted to HEK4 receptor, but that HEK4 BP may also bind to and activate other EPH-like receptor family members. As described in Example 7, HEK4 binding protein has been shown to activate both HEK4 and ECK receptors.

HEK4 binding proteins of the invention are preferably characterized by being the product of procaryotic or eucaryotic expression of an exogenous DNA sequence, i.e., HEK4 binding protein is a recombinant protein. Exogenous DNA is DNA which encodes HEK4 binding protein and includes cDNA, genomic DNA and synthetic (manufactured) DNA. HEK4 binding protein may be expressed in bacterial, yeast, plant, insect or mammalian cells in culture or in transgenic animals using DNA expression vectors appropriate for the given host cell. Expression of recombinant HEK4 binding protein in CHO cells is described in Example 3 of the specification.

Also provided is HEK4 BP in dimeric or higher order oligomeric states wherein multimeric

- 10 -

HEK4 BP is capable of binding and/or activating EPH-like receptors. Soluble HEK4 BP multimers are selected from the group consisting of HEK4 BP/immunoglobulin chimeras, HEK4 BP clustered by treatment with anti-HEK4 BP antibodies, and covalently and noncovalently attached HEK4 BP monomers. Clustered HEK4 BP is described in Example 7B and HEK4 BP chimeras are constructed using standard recombinant DNA techniques. Covalently and noncovalently attached HEK4 BP monomers are produced using protein crosslinking reagents and procedures readily available to one skilled in the art.

The polypeptides of the present invention include biologically active fragments and analogs of HEK4 BP. HEK4 BP fragments encompass amino acid sequences having truncations of one or more amino acids from the sequence shown in Figure 3 or in SEQ ID NO: 1, wherein the truncation may originate from the amino terminus, carboxy terminus, or from the interior of the protein. Analogues of the invention involve an insertion or a substitution of one or more amino acids within the sequence as shown in Figure 3 or SEQ ID NO: 1. Fragments and analogs will have at least one biological property of HEK4 binding protein, typically the ability to bind at least one EPH-like receptor.

Also encompassed by the invention are chimeric polypeptides comprising HEK4 BP amino acid sequences fused to heterologous amino acid sequences. Said heterologous sequences encompass those which, when formed into a chimera with HEK4 BP, retain one or more biological or immunological properties of HEK4 BP. In one embodiment, a HEK4 BP/immunoglobulin chimeric protein is encompassed wherein chimeric molecules may aggregate to multimeric forms of HEK4 BP for receptor binding and activation. One example is a chimera of HEK4 BP and the Fc region of IgG.

- 11 -

Also provided by the invention is an isolated nucleic acid encoding HEK4 binding protein. The nucleic acid is selected from the group consisting of:

a) the nucleic acid as shown in Figure 3 (SEQ ID NO: 1);

b) nucleic acids which hybridize under conditions of 6XSSC and 65°C with the coding regions as shown in Figure 3 (SEQ ID NO: 1);

c) nucleic acids which are degenerate to the nucleic acids of (a) and (b).

The nucleic acids may be cDNA, genomic DNA or synthetic (manufactured) DNA. It is understood that the hybridization conditions specified herein allow one skilled in the art to estimate the extent of mismatch between a given nucleic acid and a nucleic acid comprising the coding region as shown in Figure 3 (SEQ ID NO: 1) and that such conditions may be varied by changing salt, temperature and/or length of incubation or adding organic solvent at either the washing or hybridization steps and still allow one to obtain an equivalent level of mismatch during hybridization. Therefore, it is envisioned that the nucleic acids of the invention include those which hybridize with the coding regions in Figure 3 under conditions equivalent to those of 6XSSC and 65°C. Nucleic acid sequences encoding HEK 4 binding protein may have an amino terminal leader sequence and a carboxy terminal membrane anchor sequence or alternatively, may have one or both sequences removed. The encoded polypeptides will have at least one of the biological properties of HEK4 BP.

The nucleic acids of the invention will be operatively linked with nucleic acid sequences so as

- 12 -

to express HEK4 binding protein. Sequences required for expression are known to those skilled in the art and include promoters and enhancer sequences for initiation of transcription, transcription termination sites, ribosome binding sites, and sequences directing polypeptide secretion. A general description of nucleic acid sequences which serve to direct expression of exogenous genes is found in Methods in Enzymology v. 185, D.V. Goeddel, ed. Academic Press, Inc. New York (1990). Sequences directing expression of HEK4 binding protein may be homologous or heterologous. A variety of expression vectors may be used to express HEK4 binding protein in procaryotic or eucaryotic cells in culture. One such vector is pDSR $\alpha$  described in PCT Application No. WO90/14363 which was used to express HEK4 BP in CHO cells (see Example 3). In addition, vectors for tissue-specific expression of HEK4 binding protein in transgenic animals and viral-based gene transfer vectors for expression of HEK4 binding protein in human cells in vivo are also available. The nucleic acid coding regions of HEK4 binding protein may be modified by substitution of preferred codons for optimal expression in an appropriate host cell using procedures available to the skilled worker.

Plasmid pDSR $\alpha$  containing the nucleic acid sequence encoding HEK4 BP as shown in Figure 3 has been deposited with the American Type Culture Collection, Rockville, MD on \_\_\_\_\_, under ATCC Accession No. \_\_\_\_\_.

A host cell transformed or transfected with nucleic acids encoding HEK4 binding protein are also encompassed by the invention. Any host cell which produces a polypeptide having at least one of the biological properties of a HEK4 BP may be used.

- 13 -

Specific examples include bacterial, yeast, plant, insect or mammalian cells. In addition, HEK4 binding protein may be produced in transgenic animals. Transformed or transfected host cells and transgenic animals are obtained using materials and methods that are routinely available to one skilled in the art. Host cells may contain nucleic acid sequences having the full-length gene for HEK4 binding protein including a leader sequence and a C-terminal membrane anchor sequence (as shown in Figure 3) or, alternatively, may contain nucleic acid sequences lacking one or both of the leader sequence and the C-terminal membrane anchor sequence. In addition, nucleic acid fragments, variants and analogs which encode a polypeptide capable of binding HEK4 receptor may also be resident in host expression systems. Polypeptides of the invention are produced by growing transformed or transfused host cells under suitable nutrient conditions to express HEK4 BP and isolating the results at polypeptides.

Antibodies specifically binding HEK4 binding proteins of the invention are also encompassed. The antibodies can be produced by immunization with full-length (unprocessed) HEK4 binding protein or its mature forms or a fragment thereof. Antibodies may be polyclonal or monoclonal and may be human or murine-derived. Antibodies of the invention may also be recombinant, such as chimeric antibodies having the murine constant regions on the light and heavy chains replaced by human constant region sequences; or complementary determining region (CDR)-grafted antibodies wherein only the CDR is of murine origin and the remainder of the antibody chain has been replaced by human sequences.



- 14 -

The invention also provides for a pharmaceutical composition comprising a therapeutically effective amount of HEK4 binding protein and a pharmaceutically acceptable adjunct.

5 Examples of pharmaceutically acceptable adjuncts include diluents (Tris, acetate or phosphate buffers), carriers (human serum albumin), solubilizers (Tween, polysorbate), preservatives (thimerosal, benzyl alcohol) and anti-oxidants (ascorbic acid). A more

10 extensive survey of components typically found in pharmaceutical compositions appears in Remington's Pharmaceutical Sciences 18th ed. A.R. Gennaro, ed. Mack, Easton, PA (1990). As used herein, the term

15 "therapeutically effective amount" refers to that amount of HEK4 binding protein which provides a therapeutic effect for a given condition and administration regimen. Said therapeutically effective amount may vary from 0.01 µg/kg body weight

20 to 10 mg/kg body weight and may be determined by one skilled in the art.

HEK4 binding protein may be administered by injection, either subcutaneous, intravenous or intramuscular, or by oral or nasal administration.

25 The route of administration to be chosen will depend upon several variables, including the nature and severity of the condition being treated and the pharmacokinetic properties of the HEK4 binding protein preparation. HEK4 binding protein may be formulated

30 for delivery in a particular fashion, e.g., it may be modified with water soluble polymers, such as polythylene glycol to improve properties for nasal delivery or to improve serum half-life after injection; or it may be incorporated into particulate

35 preparations of polymeric compounds (e.g., liposomes)

- 15 -

for controlled delivery over an extended period of time.

The expression of HEK4 receptor and HEK4 BP  
5 in various tissues is reported in Examples 6A and 6B,  
respectively. HEK4 receptor mRNA was most abundant in  
human placenta and was also detected in heart, brain,  
lung, liver, muscle, kidney tissues. HEK4 BP mRNA was  
most abundant in human adult brain, kidney and  
10 placenta, and was detected at lower levels in heart,  
lung, liver, spleen, prostate, testis, ovary, small  
intestine, muscle, pancreas and colon. These patterns  
of expression suggest that activation of HEK4 receptor  
by HEK4 BP modulates the growth and/or differentiation  
15 of a variety of target cells, particularly those in  
the brain, heart, lung, liver, muscle and pancreas  
where expression of both receptor and ligand are  
detected. In addition, Wicks et al., supra has  
reported HEK4 receptor mRNA in pre-B and T cell lines,  
20 suggesting a role for HEK4 BP in hematopoiesis.

As described in Example 7, HEK4 BP also  
activates ECK receptor in a cell-cell  
autophosphorylation assay. Eck receptor mRNA is most  
abundant in adult rat lung, small intestine, kidney,  
25 ovary and skin with lower levels detected in brain,  
spleen and submaxillary gland (Lindberg and Hunter,  
supra). Recently, it has been shown that Eck is  
expressed in the nervous system of the early mouse  
embryo (Becker et al. Mech. Dev. 47, 3-17 (1994);  
30 Ganju et al. Oncogene 9, 1613-1624 (1994)). These  
observations suggest that activation of ECK receptor  
by HEK4 BP may modulate the growth and/or  
differentiation of cells expressing ECK, such as those  
in the lung, intestine, kidney, skin and nervous  
35 system.

- 16 -

Therefore, HEK4 BP is useful in modulating (i.e., increasing or decreasing) the extent of growth and/or differentiation of target cells in various tissues. The target cells will have at least one  
5 receptor which is activated by HEK4 BP wherein the receptor is preferably a member of the EPH sub-family of receptor PTKs. Potential therapeutic uses for HEK4 BP are described below.

One aspect of the invention is the use of  
10 HEK4 BP to modulate cell growth and differentiation in the nervous system. HEK4 BP may be used to maintain or restore cellular function in the nervous system of a mammal which has been decreased or eliminated by disease or injury or is at risk of being decreased or  
15 eliminated by disease or injury. Target cells include neurons and glial cells. Conditions that may be treated by HEK4 BP include central nervous system disorders such as Alzheimer's disease, Parkinson's disease, multiple sclerosis, stroke and Huntington's  
20 disease and peripheral nervous system disorders such as amyotrophic lateral sclerosis (ALS) and peripheral neuropathies. Physical injuries to the spinal cord and to peripheral neurons may also be treated with HEK4 BP.

25 Another aspect of the invention is the modulation by HEK4 BP of growth and differentiation of digestive tract (including large and small intestine), liver, lung, pancreas, muscle and hematopoietic tissues. This activity of HEK4 BP may be particularly  
30 useful in regeneration of tissue in these and other sources which has been damaged or depleted by disease or injury.

It has been observed that the sub-family of EPH-like receptors and their corresponding ligands are  
35 highly expressed in some carcinoma cell lines (see for example, PCT Application No. WO94/11020 for expression

- 17 -

of ECK receptor and ECK binding protein in human carcinoma cell lines). Thus another aspect of the invention is the treatment of cancers using HEK4 BP antagonists to block cell proliferation. Such cancers  
5 are likely to be associated with organs which express HEK4 receptors and/or Eck receptors. HEK4 BP antagonists may be any compound which blocks the biological activity of HEK4 BP and may include, but are not limited to, the following: antibodies which  
10 bind to either HEK4 BP or to an EPH sub-family receptor which is activated by HEK4 BP such that a receptor/ligand interaction is prevented; HEK4 BP which binds to, but does not activate and EPH-like receptor; and soluble EPH-like receptors which bind to  
15 HEK4 BP. It is envisioned that small molecule mimetics of the above described antagonists are also encompassed by the invention.

In addition to in vivo applications, HEK4 BP may also be used ex vivo to amplify cell  
20 populations prior to transplantation. It is envisioned that HEK4 BP may promote growth in culture of cells from the digestive tract, liver, lung, bone marrow, kidney, or central and peripheral nervous systems (and glial cells) neurons such that the  
25 amplified population can be introduced back into a patient in need of such therapy. Such so-called "cell therapy" is useful in replenishing cells after damage or depletion and may be appropriate under conditions where systemic administration of HEK4 BP is not  
30 preferred.

HEK4 BP may be used alone or in combination with other therapeutic agents for the treatment of cancer, neurological disorders, disorders of the digestive tract, liver, or lung, and for the ex vivo  
35 expansion of cell populations. HEK4 BP may be used in conjunction with other chemotherapeutic drugs or with

- 18 -

radiation therapy for the treatment of cancer, or with other neurotrophic factors such as brain derived neurotrophic factor (BDNF), ciliary neurotrophic factor (CNTF), neurotrophin-3 (NT-3), nerve growth factor (NGF), or glial derived neurotrophic factor (GDNF) for neurological disorders; or with tissue growth factors such as platelet derived growth factor (PDGF), fibroblast growth factor (FGF), epidermal growth factor (EGF), hepatocyte growth factor (HGF) or keratinocyte growth factor (KGF) for restoration of damaged or depleted tissues.

Isolated nucleic acids of the present invention are useful reagents for the detection and quantitation of DNA and/or RNA coding for HEK4 BP by standard hybridization procedures such as those described in Sambrook et al. Molecular Cloning. A Laboratory Manual, 2d ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989). These reagents allow one to determine the potential of various cell types to express HEK4 BP and related polypeptides and are also useful for detecting abnormalities in genes encoding HEK4 BP or in sequences controlling the expression of HEK4 BP. Nucleic acids of the invention are also useful for controlling expression levels of HEK4 BP. So-called "anti-sense" nucleic acids hybridize to DNA and/or RNA strands encoding HEK4 BP in a manner that blocks transcription or translation of HEK4 BP nucleic acid sequences. Introduction of HEK4 BP anti-sense nucleic acids into cells overexpressing HEK4 BP is appropriate when such overexpression leads to undesirable physiological effects, such as excessive cell proliferation.

Antibodies which specifically bind HEK4 BP are useful reagents for the detection and quantitation

- 19 -

of HEK4 BP in biological samples using immunoassays (Western blots, RIAs, ELISAs) that are conventional to the art. The presence of HEK4 BP may be indicative of cell proliferation or the potential for cell proliferation, and elevated levels may signal abnormal cell growth typically associated with cancer. In addition, antibodies of the invention may also be useful therapeutic reagents that act as agonists or antagonists of HEK4 BP activity. Antibodies may bind to HEK4 BP in a manner that directly or indirectly blocks HEK4 BP binding an EPH receptor (either HEK4 or ECK). Alternatively, antibodies may bind to HEK4 BP in a manner which promote receptor binding and activation by, for example, "clustering" HEK4 BP into a dimeric or higher multimeric forms to allow more efficient binding and activation of receptor. Antibodies can be monoclonal, polyclonal, or recombinant.

The following examples are offered to more fully illustrate the invention, but are not construed as limiting the scope thereof.

#### EXAMPLE 1

##### Production of HEK4 receptor extracellular domain (HEK4-X)

A cDNA clone coding for the HEK4 receptor protein tyrosine kinase was isolated from a human fetal brain cDNA library (Stratagene, La Jolla, CA) as described in co-pending and commonly owned U.S. Serial No. 08/229,509. The sequence of this clone was identical to that in Figure 1 of Wicks et al., supra with the following exceptions. Wicks reported the

- 20 -

sequence TTA at nucleotides 1618-1620 whereas the HEK4  
receptor clone isolated as described here had the  
sequence TTC at these positions. However, Wicks et  
al.'s predicted protein sequence specifies a  
5 phenylalanine residue in this position, which is  
inconsistent with an "A" at nucleotide 1620 (TTA codes  
for leucine while TTC codes for phenylalanine). Also,  
nucleotides 1529 through 1531 of the Wicks et al.  
sequence are absent from the sequence obtained here.  
10 This change does not affect the translational reading  
frame, but does eliminate the predicted glutamine  
residue at position 478 of the Wicks sequence. The  
effect of these differences on the biological activity  
of the receptor or the ability to bind ligand is  
15 unknown.

The HEK4 receptor cDNA clone was used as a  
template in a polymerase chain reaction (PCR) designed  
to amplify a DNA fragment coding for the ligand  
binding domain of the HEK4 receptor. The primers used  
20 were:

433-26) 5' GGATCTAGAGCACCAGCAACATGGATTGT 3'  
(SEQ ID NO: 3)

25 409-10) 5' TCGGTCTAGATCATTATTGGCTACTTTCACCAGAGAT 3'  
(SEQ ID NO: 4)

These primers produce a fragment 1656  
nucleotides in length that codes for a protein of 540  
30 amino acids. The predicted protein consists of the  
entire extracellular domain of the HEK4 receptor from  
the amino terminus up to but not including the  
transmembrane region. The 1656 nucleotide fragment  
was digested with the restriction endonuclease XbaI  
35 and ligated into the expression vector pDSR $\alpha$  which had  
been digested with the same enzyme. The resulting

- 21 -

expression plasmid was introduced into CHO cells by calcium phosphate mediated transfection (Cellfect, Pharmacia, Piscataway, NJ). Individual colonies were selected based upon the expression of the

5 dihydrofolate reductase (DHFR) gene which resides on the expression plasmid. Expression of the HEK4 gene was monitored by RNA solution hybridization (Hunt et al. Exp. Hematol. 19, 779-784 (1991)) and/or by

10 Western blotting with antibodies directed against amino acids 22-148 of the HEK4 extracellular domain.

HEK4 expression was enhanced by growth of the selected clones in 100 nM methotrexate. One of the pDSR $\alpha$ /HEK4-X clones was chosen for large scale production. Twenty-four roller bottles were seeded at

15 a density of approximately  $2 \times 10^7$  cells/bottle in 200 ml each of Dulbecco's Minimal Essential Media (DMEM) supplemented with non-essential amino acids (1X NEAA, Gibco), 100 nM methotrexate, 1X

20 penicillin/streptomycin/glutamine (1X PSG, Gibco) and 10% fetal bovine serum. Cells reached confluence in 3-4 days at which time the media was changed to DMEM/NEAA/PSG lacking serum. Cell-conditioned media was harvested after seven days, concentrated, and diafiltered against 10 mM Tris-HCl, pH 8.5. The

25 concentrated media was loaded onto a Q-sepharose FF (Pharmacia) ion exchange column and bound material was eluted with a linear gradient of 0 to 0.5 M NaCl in 10 mM Tris-HCl, pH 8.5. Fractions were analyzed by

30 SDS-PAGE and western blotting using a rabbit polyclonal antibody directed against residues 22-148 of the HEK4 external domain. Fractions containing HEK4-X protein were pooled, concentrated and loaded onto an S-200 (Sephacryl S-200, Pharmacia) column. Fractions from this column were analyzed as before and

35 those containing HEK4-X were pooled.



## EXAMPLE 2

A. Purification of HEK4-X Binding Activity

5 We have previously described the use of the BIAcore™ instrument (Pharmacia Biosensor, Piscataway, NJ) for the detection of receptor binding activity in concentrated cell-conditioned media (Bartley et al., supra). We used a similar strategy to screen for HEK4  
10 receptor binding activity as described below.

The surface of a BIAcore sensor chip was activated by injection of 0.2 M 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide-HCl and 0.05 M N-hydroxysuccinimide at a flow rate of 5 ul/min.  
15 Purified HEK4-X at a concentration of 250 ug/ml was applied to the activated surface in two 50 ul injections at the same flow rate. Unreacted binding sites were blocked by injection of 1M ethanolamine, pH 8.5. The surface was washed overnight in 10 mM  
20 HEPES, 150 mM NaCl, 3.4 mM EDTA, 0.005% Tween 20, pH 7.4 until the baseline was stable. Typically, immobilization resulted in 6000-8000 resonance units (RU) of HEK4-X bound to the sensor chip.

Conditioned media samples were collected  
25 from 108 cell lines grown either without fetal bovine serum (FBS) or in the presence of 0.5% FBS. Conditioned media produced under serum-free conditions was adjusted to 0.5% FBS before further processing. The media was filtered, concentrated 25-fold, and  
30 stored in aliquots at -80°C. 30ul samples of each medium were injected onto the HEK4-X surface at a flow rate of 5 µl/min and the binding response measured 20 seconds after the conclusion of each injection. Between samples, the surface was regenerated by  
35 10-15 µl injections of 25 mM 3-(cyclohexylamino)-1-propanesulfonic acid, pH 10.4.

- 23 -

Concentrated conditioned media samples displaying binding of 200 resonance units or more are listed in Table 1.

5

Table 1

<u>Cell Line</u>	<u>Description</u>	<u>Binding</u> <u>(resonance units)</u>
10		
	HCT116 human colon carcinoma	911
	M-14 human melanoma	337
	LS174T human colon adenocarcinoma	316
	A498 human kidney carcinoma	274
15	A172 human glioblastoma	269
	PK(15)1 porcine kidney	234
	JEG-1 human choriocarcinoma	220
	Y-79 human retinoblastoma	216
	HT 1080 human fibrosarcoma	200
20		

The five conditioned media displaying the most binding were selected for further investigation. When soluble dextran was added to samples before  
25 injection to reduce non-specific binding, the signals from HCT116 and LS174T conditioned media were greatly reduced. A172 cells proved difficult to grow and were unsuitable for the large scale production of conditioned media. Based on these experiments and on  
30 pilot scale receptor affinity chromatography, the A498 cell line (ATCC No. HTB 44) was chosen as the best source of conditioned media for purification of a HEK4 binding protein.

A HEK4 receptor affinity column was  
35 prepared by immobilization of HEK4-X on CNBr-activated Sepharose 4B (Pharmacia). Purified HEK4-X was dialyzed against 0.1 M NaHCO<sub>3</sub>, 0.5 M NaCl, pH 8.3 and

- 24 -

brought to a final concentration of 2 mg/mL.

Immobilization of HEK4-X was done at a ligand density of 1 mg/mL according to the method of Kenny et al.

(New Protein Techniques, J.M. Walker, ed. The Humana

5 Press, Clifton, NJ 1988). Forty liters of A498

conditioned media produced in 0.5% serum-containing

media was concentrated 40-fold, diafiltered against

PBS and 0.02% NaN<sub>3</sub>, and loaded onto the HEK4-CNBr

sepharose column. The column was washed with PBS and

10 bound material was eluted with 50 mM sodium acetate,

0.5 M sodium chloride, pH 4.0. Fractions were

collected in 1 mM CHAPS and loaded directly on

polyacrylamide gels. Gels were either stained with

silver for analysis or blotted onto PVDF membranes

15 (Problot, Applied Biosystems, Foster City, CA) in

preparation for N-terminal amino acid sequencing

(Fausset & Lu, 1991).

The pH 4.0 elution fractions from the

HEK4-CNBr Sepharose column contained three major

20 protein species with molecular weights of 21, 25 and

27 kD which were not apparent in the load or wash

fractions. The fractions containing these 3 proteins

were pooled, concentrated in the presence of CHAPS and

applied to a Vydac C4 reverse phase HPLC column (4.6 x

25 150 mm). The column was eluted with a gradient of

acetonitrile (26-35%) in 0.1% trifluoroacetic acid.

Fractions were collected, volume reduced under vacuum,

and analyzed for HEK4 binding protein. The three

major peaks detected by absorbance at 214 nm were

30 pooled and analyzed by SDS-PAGE (see Figure 2).

Further purification of the three isoforms of HEK4 BP

was achieved by reapplying the proteins to the same C4

column.

- 25 -

**B. Sequencing of peptides**

A sample from the initial purification of A498 cell-conditioned media was submitted for protein sequencing. The sample was analyzed on SDS-PAGE and  
5 blotted onto PVDF membrane. The gel band identified as the HEK4 binding protein was excised and analyzed for five cycles on an Applied Biosystems 477A protein sequencer. This yielded no sequence, indicating that the protein was N-terminally blocked. The sample was  
10 then treated with cyanogen bromide and reapplied to the sequencer. A tentative sequence was obtained from the cleaved sample by assuming the highest yield at each cycle to belong to the same peptide. Even given this assumption, recoveries were so small as to render  
15 the sequence unreliable after cycle 10. The sequence obtained in this manner is shown as peptide #1 in Table 2.

20

**Table 2**

<u>Peptide No.</u>	<u>Amino acid sequence</u>
1	Val-Asn-Phe-Asp-Gly-Tyr-Ser-Ala-Arg- 25 Asp (SEQ ID NO: 5)
2	Val-Phe-Asp-Val-Asn-Phe-Lys-Val-Glu-X- Ser-Leu-Glu-Pro-Ala-Asp (SEQ ID NO: 6)
3	Ala-Val-Ala-Asp-Arg-Tyr-Ala-Val-Tyr- 30 Trp-Asn-Ser-Ser-Asn-Pro-Arg-Phe-Gln- Arg-Gly-Asp-Tyr-His-Ile-Ile-Val-X-Ile- Asn-X-Tyr (SEQ ID NO: 7)

35

- 26 -

Subsequent analysis of samples cleaved by cyanogen bromide, then separated by SDS-PAGE indicated that position 9 of peptide #1 was a cysteine residue and position 6 of peptide #2 was aspartic acid.

5 Position 25 of peptide #3 was subsequently found by DNA sequencing to be aspartic acid. The sequence data shown in peptides #2 and #3 in Table 2 was obtained by analysis of tryptic digests of the protein followed by separation of the resulting peptides on a microbore C4  
10 column. These experiments were done with larger amounts of starting material and therefore yielded more reliable sequence and allowed 20-30 cycle sequencing runs. Comparison of the peptide sequences in Table 2 with B61 suggests that they represent  
15 fragments of a related protein. We therefore conclude that the HEK4 binding protein is another ligand for the human EPH-like kinase sub-family.

20

## EXAMPLE 3

A. Cloning and Sequencing of cDNA Encoding HEK4 Binding Protein

The amino acid sequences obtained from HEK4  
25 BP peptides as shown in Table 2 were used to design oligonucleotide primers. Primers 702-3 and 633-11 were used in a PCR reaction with random primed A498 cDNA as a template.

30 702-3) 5' GAYMGNTAYGCNGTNTAYTGG 3' (SEQ ID NO:8)

633-11) 5' RTANCCRTCRAARTTNACCAT 3' (SEQ ID NO:9)

The 175 base pair fragment amplified by these primers  
35 was sequenced and found to be closely related to B61. This fragment was then radiolabeled with <sup>32</sup>P by random

- 27 -

priming and used as a probe to screen a cDNA library for clones containing the full length HEK4 BP cDNA. An oligo-dT primed human placental cDNA library purchased from Stratagene (La Jolla, CA) was plated at a density of 30,000 plaques/150 mm plate. Replicas of the plaques arrayed on the plates were made on GeneScreen<sup>TM</sup> hybridization transfer membranes (New England Nuclear, Boston, MA) as directed by the manufacturer. Two replica filters were made for each plate. Filters were pre-hybridized in 6X SSC, 1X Denhardt's buffer, 50 ug/ml salmon testis DNA, 1% SDS at 65°C for 4 hours followed by hybridization with the <sup>32</sup>P-labeled probe for 12 hours under the same conditions. Following hybridization, filters were washed two times, 1 hour each, in 0.2X SSC, 0.5% SDS at 65°C and exposed to Kodak XAR film overnight with an intensifying screen. Comparison of the two filters made from each plate showed that five plaques were positive on both replicas. The phage around each positive plaque were removed, resuspended in buffer, and replated at a lower density to produce well-separated plaques for secondary screening. Individual plaques which were positive upon rescreening (using the same method as the primary screen) were picked. The inserts from these phage were transferred into the pBluescript plasmid by in vivo excision as described by the manufacturer (Stratagene). Three of the five inserts were identical and contained the entire coding region of HEK4 BP while the other two represented overlapping clones. A consensus sequence was assembled using data from the three inserts containing the entire coding region and is shown in Figure 3.

The HEK4 BP cDNA sequence predicts a protein of between 213 and 228 amino acids, depending on which of three possible initiator codons is utilized. Based upon rules for translation of

- 28 -

vertebrate mRNAs (Kozak Cell 44, 283-292 (1986)), the third in-frame ATG is an unlikely initiator, while the first ATG, being the farthest upstream is the most likely to be the principal initiation codon. As for

5 B61, HEK4 BP has hydrophobic amino acids on both the amino and carboxy termini. These probably function as a secretion signal sequence and a membrane anchor, respectively. Like B61, HEK4 BP apparently has both soluble and membrane bound forms. Although we were

10 not able to obtain N-terminal protein sequence data, we would predict cleavage of the signal peptide to yield a mature protein with serine at position 1 (Figure 3). Based on peptide mapping and mass spectrometric analysis, proline-179 (Figure 3) appears

15 to be the C-terminal amino acid in the major soluble form found in A498 cell-conditioned media. An alternate form with alanine-177 (Figure 3) at the C-terminus was also detected.

20 B. Expression of recombinant HEK4 BP

The HEK4 BP cDNA clone shown in Figure 3 was inserted into the plasmid vector pDSR $\alpha$  for expression mammalian cells. The recombinant plasmid was transfected into Chinese hamster ovary (CHO) cells

25 by calcium phosphate precipitation and cells containing the plasmid were selected by growth in DMEM (high glucose, GIBCO, Bethesda, MD), 1X penicillin/streptomycin/glutamine (PSG), 1X non-essential amino acids (NEAA) containing 10% fetal

30 bovine serum (FBS), but lacking HT supplement (HT supplement: 10 mM sodium hypoxanthine, 1.6 mM thymidine). The expression of HEK4 BP in several clones was assessed by the level of HEK4-receptor binding activity in each clone's cell-conditioned

35 media as determined by BIAcore (Pharmacia Biosensor, Piscataway, NJ). This correlated well with the level

- 29 -

of HEK4 BP mRNA in the clones as determined by Northern blot hybridization. One clone, CHO/HL6, was a significantly better producer of recombinant HEK4 BP than the others and was chosen for further work.

5 Expression of HEK4 BP by CHO/HL6 was enhanced 2 to 4-fold by treatment with increasing amounts of methotrexate up to 100 nM over a period of several weeks. Following amplification, cells expressing HEK4 BP were expanded and transferred to roller

10 bottles for production of conditioned media to be used as a source for purification of the recombinant protein. A total of 100 roller bottles were seeded with CHO/HL6 cells at  $10^7$  cells per bottle. Cells were grown until confluent (approximately 4 days) in

15 DMEM (high glucose, GIBCO, Bethesda, MD) , 1X PSG, 1X NEAA, and 10% FBS. Following the growth phase, the media was removed replaced with the same media but with 0.5 % rather than 10% FBS. After 3 days, the media was collected, filtered to remove any cell

20 debris, and stored frozen at  $-80^{\circ}\text{C}$ .

## EXAMPLE 4

25 Purification of recombinant HEK4 BP

Approximately 20-25 liters of conditioned medium from CHO/HL6 cells were thawed at room temperature and filtered. The medium was concentrated and diafiltered

30 against 10mM Tris-HCl, pH 8.5 ( $4^{\circ}\text{C}$ ) using a 10,000 molecular weight cut off membrane. The diafiltrate was applied to a column of Q-Sepharose, High Performance and subsequently eluted with a linear gradient of NaCl (0-0.3 M) in 10 mM Tris-HCl, pH 8.5.

35 Fractions were analyzed for the presence of HEK4 BP by immunoblotting using an antibody generated against



- 30 -

unfolded HEK4 BP produced in *E. coli*, or by binding to HEK4 receptor immobilized on a BIAcore sensor chip. Fractions containing HEK4 BP were pooled, concentrated and applied to a gel filtration column. (Superdex 75, 5 x 85 cm, PBS, 3 mL/min). Fractions containing HEK4 BP were further purified by C4 reverse phase HPLC (Vydac 214TP 4.6 x 250 mm, 2.9 mL/min) using an acetonitrile gradient (22-44%) in 0.1% trifluoroacetic acid. The column profile and SDS-PAGE analysis of peak fractions are shown in Figure 4. Fractions were evaporated under vacuum and formulated in 0.25 M Tris-HCl, 2 mM CHAPS, pH 7.5.

15

## EXAMPLE 5

A. Production of Antibodies to HEK4 Receptor

Antibodies directed against the HEK4 receptor extracellular domain were produced using standard methods (Harlow and Lane, Antibodies: A Laboratory Manual Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1988)). cDNA encoding amino acids 22-148 of the HEK4 receptor was inserted into the pATH vector (Yansura, 1990) in the same translational reading frame as the TrpE gene. The resulting plasmid was introduced into an *E. coli* host strain resulting in expression of the HEK4/TrpE fusion protein. Bacterial cell lysates were fractionated by preparative SDS-PAGE and the band containing the HEK4/TrpE fusion was excised. The crushed gel slice was used to immunize rabbits according to standard protocols (Harlow and Lane, supra). The antisera generated by this method recognized both the HEK4/TrpE antigen and recombinant HEK4-X produced in CHO cells.

Antibodies to the C-terminal 12 amino acids of the HEK4 receptor (sequence is cys-leu-glu-thr-gln-

- 31 -

ser-lys-asn-gly-pro-val-pro-val) were produced by the same method (Harlow and Lane, supra) using a synthetic peptide chemically linked to keyhole limpet hemocyanin (KLH) as the antigen. The antiserum was purified by,  
5 passage over a column upon which peptide antigen had been immobilized using a SulfoLink kit (Pierce, Rockford, IL). These antibodies were able to specifically recognize HEK4 receptor by Western blots.

10 B. Production of Antibodies to HEK4 BP

HEK4 BP cDNA as shown in Figure 3 was used as a template for PCR with primers 819-31 and 819-28 to produce a polypeptide fragment coding for amino acids 1-179 of HEK4 BP (Figure 3).

15

819-31) 5' GGAGGACATATGAGCCAGGACCCGGGCTCCAAG 3'  
(SEQ ID NO:10)

20

819-28) 5' GAAGAAGGATCCCTATGGCTCGGCTGACTCATGTAC 3'  
(SEQ ID NO:11)

The PCR fragment was cloned into the expression vector pCFM1656 using the NdeI and BamHI sites included in the primers. The resulting recombinant plasmid was  
25 transformed into E. coli FM5 (ATCC No. 53911) and the truncated HEK4 BP was expressed as insoluble inclusion bodies. The inclusion bodies were solubilized and the HEK4 BP fragment purified by SDS-polyacrylamide gel electrophoresis was used as an antigen in rabbits.  
30 The antisera was generated and characterized as described (Harlow and Lane, supra) and recognized HEK4 BP expressed in CHO cells by Western blot analysis.

- 32 -

## EXAMPLE 6

A. Expression pattern of the HEK4 receptor

5           The expression of HEK4 receptor mRNA in rat  
and human tissues has been previously reported in co-  
pending and commonly owned U.S. Serial No. 08/229,509,  
relevant portions of which have been incorporated  
herein by reference. The results of these studies are  
10 summarized in Table 3.

TABLE 3

Tissue Distribution of HEK4 Receptor

<u>Tissue:</u>	<u>Human</u>	<u>Rat</u>
Brain	++	+
Heart	+	bd
Kidney	+	bd
Liver	+	bd
Lung	+	+
Muscle	+	bd
Ovary	nt	bd
Pancreas	+	nt
Placenta	+++	nt
Stomach	nt	bd
Testis	nt	+
Thymus	nt	bd

15

bd = below detection  
nt = not tested

20 In the human tissues studied, HEK4 receptor mRNA is  
most abundantly expressed in placenta, with lower  
levels in heart, brain, lung, and liver. Previous  
studies on HEK4 receptor mRNA in cell lines found

- 33 -

expression in one pre-B cell line and two T-cell lines (Wicks et al. 1992).

B. Expression pattern of HEK4 BP

5           The expression of HEK4 BP mRNA in human tissues was examined by Northern blot analysis. A Northern blot containing 2 µg of polyA<sup>+</sup> from each of the tissues indicated was purchased from Clontech (Palo Alto, CA) and hybridized with a <sup>32</sup>P-labeled  
10 HEK4 BP probe. As shown in Figure 5, HEK4 BP mRNA is expressed at high levels in human adult brain, kidney, and placenta. Readily detectable levels can also be found in heart, lung, liver, spleen, prostate, testis, ovary, small intestine, and colon. The presence of  
15 HEK4 BP mRNA in many different tissues is consistent with the idea that this factor is important for the differentiation, development, and/or maintenance of a variety of cell types.

20

EXAMPLE 7

A. HEK4 BP activation of EPH subfamily receptors by cell-cell autophosphorylation

25           The hallmark of receptor activation for all known receptor protein-tyrosine kinases is autophosphorylation (van der Geer et al., *supra*). To determine whether HEK4 BP can activate HEK4 receptor, a cell-cell autophosphorylation assay was performed.  
30 Recipient cells were CHO cells transfected with HEK4 receptor cDNA which had been serum-starved by incubation in media with 0.5% serum for 16 hours. The donor cells were CHO cells transfected with HEK4 BP cDNA (see Example 3B) or CHO cells that had been  
35 transfected with vector alone. Donor cells were scraped from the surface of their growth vessel in

- 34 -

phosphate-buffered saline and added to recipient cells for 30 minutes at 37°C. After washing, the recipient cells were lysed in modified RIPA buffer (10 mM sodium phosphate, pH 7.4, 150 mM sodium chloride, 0.1% sodium dodecyl sulfate, 1% NP-40, 1% deoxycholate, 10 mg/ml aprotinin, 5mM EDTA, 200 mM sodium orthovanadate). Receptors were immunoprecipitated from the cell lysate and prepared for SDS polyacrylamide gel electrophoresis as previously described (Bartley et al., supra). After electrophoresis and electroblotting to membranes, the immunoprecipitates were probed with antiphosphotyrosine antibodies (4G10, UBI, Lake Placid, NY). Immune complexes were detected by horseradish peroxidase conjugated secondary reagents using chemiluminescence as described by the manufacturer (ECL, Amersham). As shown in Figure 6, cells expressing HEK4 binding protein were able to stimulate tyrosine phosphorylation on both the HEK and ECK receptors. Control cells did not stimulate the phosphorylation of either receptor. The results demonstrate that HEK4 BP can activate both the HEK4 and the ECK receptors.

#### B. Activation of HEK4 Receptor by Soluble HEK4 BP

To determine whether soluble recombinant HEK4 BP could activate the HEK4 receptor, CHO cells transfected with HEK4 receptor cDNA were treated with conditioned media from CHO cells expressing HEK4 BP (see Example 3B) or with purified recombinant HEK4 BP (see Example 4). The cells were serum-starved by incubation in media with 0.5% serum for 16 hours prior to the treatments. Treatments were for 30 minutes at 37°C, after which the cells were lysed in modified NP40 buffer (50 mM Tris, pH 8.0, 150 mM sodium chloride, 1% NP40, 10 mg/ml aprotinin, 5mM EDTA, 200 mM sodium orthovanadate), HEK4 receptor was

- 35 -

immunoprecipitated, and prepared for SDS polyacrylamide gel electrophoresis as previously described (Bartley et al., supra). After electrophoresis and electroblotting to membranes, the immunoprecipitates were probed with antiphosphotyrosine antibodies (4G10, UBI, Lake Placid, NY). Immune complexes were detected by horseradish peroxidase conjugated secondary reagents using chemiluminescence as described by the manufacturer (ECL, Amersham). As shown in Figure 7, soluble recombinant HEK4 BP in conditioned media and after purification activated HEK4 receptor. This activation was enhanced by pretreatment of conditioned media or purified HEK4 BP with the antibodies of Example 5B which had been affinity purified on a HEK4 BP column. Antibodies (~50 µg/ml) were incubated with conditioned medium or purified HEK4 BP at 4° for 1 hr. prior to treatment of CHO cells expressing HEK4 receptor.

20

## EXAMPLE 8

Affinity of HEK4 BP for EPH-like Receptors

25

A competition assay was used to measure differences in HEK4 BP binding to different EPH-like receptors. Purified HEK4 BP was incubated with various concentrations of either HEK4, HEK8 or ECK soluble receptors and binding of the mixture to immobilized HEK4 receptor was analyzed by BIAcore. The concentration of soluble receptor that inhibited HEK4 BP binding by 50% is termed IC50. IC50 values allow a comparison of the relative affinity of HEK4 BP for related receptors.

35

- 36 -

IC50 values were determined as follows. Analysis of HEK4 BP binding to immobilized HEK4 receptor showed a linear response in the range 60 to 500 ng/ml. Various amounts of the purified extracellular domains of HEK4, ECK, or HEK8 were incubated with 0.250 µg/ml HEK4 BP prepared as described in Example 4 in solutions containing 100 µg/ml BSA, 10mM HEPES, 0.15M NaCl, 3.4 mM EDTA, and 1 mg/ml soluble dextran, pH 7.4. These solutions were incubated for at least 30 minutes at 3°C prior to injection. Protein concentrations of receptor stocks were confirmed by BCA protein assay. Duplicates of each sample were run in parallel with standard curves on two different days. All surfaces were regenerated to within 10 RU of baseline with 25mM CAPS and 1M NaCl pH 10.4. The mean binding response was plotted versus soluble receptor concentration (Figure 8) providing IC50 values of 0.55 ug/ml for HEK4, 5.0 µg/ml for ECK, and 10.5 µg/ml for HEK8. Thus, HEK4 BP preferentially binds HEK4 receptor compared to two other EPH family members, ECK and HEK8.

25

\* \* \*

While the invention has been described in what is considered to be its preferred embodiments, it is not to be limited to the disclosed embodiments, but on the contrary, is intended to cover various modifications and equivalents included within the spirit and scope of the appended claims, which scope is to be accorded the broadest interpretation so as to encompass all such modifications and equivalents.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: Amgen Inc.
- (ii) TITLE OF INVENTION: Ligands for EPH-Like Receptor
- (iii) NUMBER OF SEQUENCES: 11
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: Amgen Inc.
  - (B) STREET: 1840 Dehavilland Drive
  - (C) CITY: Thousand Oaks
  - (D) STATE: California
  - (E) COUNTRY: USA
  - (F) ZIP: 91320-1789
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.25
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE:
  - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: Winter, Robert B.
  - (C) REFERENCE/DOCKET NUMBER: A-325

## (2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 1728 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 175..858
- (ix) FEATURE:
  - (A) NAME/KEY: mat\_peptide
  - (B) LOCATION: 232..858
- (ix) FEATURE:
  - (A) NAME/KEY: sig\_peptide
  - (B) LOCATION: 175..231



- 38 -

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAATTC	CCCC	AGCTTGGTGG	GCGCCTCTTT	CCTTTCTCGC	CCCCTTTCAT	TTTTATTTAT	60
TCATATTTAT	TTGGCGCCCG	CTCTCTCTCT	GTCCCTTTGC	CTGCCTCCCT	CCCTCCGGAT		120
CCCCGCTCTC	TCCCCGGAGT	GGCGCGTCGG	GGGCTCCGCC	GCTGGCCAGG	CGTG	ATG	177
						Met	
						-19	
TTG	CAC	GTG	GAG	ATG	TTG	ACG	225
Leu	His	Val	Glu	Met	Leu	Thr	
			-15			-10	
GTG	TTC	AGC	CAG	GAC	CCG	GGC	273
Val	Phe	Ser	Gln	Asp	Pro	Gly	
		1				5	
GTC	TAC	TGG	AAC	AGC	AGC	AAC	321
Val	Tyr	Trp	Asn	Ser	Ser	Asn	
		15				20	
ATT	GAT	GTC	TGT	ATC	AAT	GAC	369
Ile	Asp	Val	Cys	Ile	Asn	Asp	
			35			40	
GAG	GAC	TCC	GTC	CCA	GAA	GAT	417
Glu	Asp	Ser	Val	Pro	Glu	Asp	
			50			55	
GTG	AAC	TTT	GAT	GGC	TAC	AGT	465
Val	Asn	Phe	Asp	Gly	Tyr	Ser	
		65				70	
AAG	AGA	TGG	GAA	TGT	AAC	CGG	513
Lys	Arg	Trp	Glu	Cys	Asn	Arg	
		80				85	
TTC	TCT	GAA	AAA	TTC	CAG	CTC	561
Phe	Ser	Glu	Lys	Phe	Gln	Leu	
		95				100	
TTC	AGG	CCA	GGC	CGA	GAA	TAT	609
Phe	Arg	Pro	Gly	Arg	Glu	Tyr	
			115			120	
AAT	GGA	AGA	AGG	TCC	TGT	CTA	657
Asn	Gly	Arg	Arg	Ser	Cys	Leu	
			130			135	
AAT	AGC	TGT	ATG	AAA	ACT	ATA	705
Asn	Ser	Cys	Met	Lys	Thr	Ile	
			145			150	

- 39 -

AAC GAC AAA GTA GAA AAT TCA TTA GAA CCA GCA GAT GAC ACC GTA CAT	753
Asn Asp Lys Val Glu Asn Ser Leu Glu Pro Ala Asp Asp Thr Val His	
160 165 170	
GAG TCA GCC GAG CCA TCC CGC GGC GAG AAC GCG GCA CAA ACA CCA AGG	801
Glu Ser Ala Glu Pro Ser Arg Gly Glu Asn Ala Ala Gln Thr Pro Arg	
175 180 185 190	
ATA CCC AGC CGC CTT TTG GCA ATC CTA CTG TTC CTC CTG GCG ATG CTT	849
Ile Pro Ser Arg Leu Leu Ala Ile Leu Leu Phe Leu Leu Ala Met Leu	
195 200 205	
TTG ACA TTA TAGCACAGTC TCCTCCCATC ACTTGTACACA GAAAACATCA	898
Leu Thr Leu	
GGGTCTTGGG ACACCAGAGA TCCACCTAAC TGCTCATCCT AAGAAGGGAC TTGTTATTGG	958
GTTTTGGCAG ATGTCAGATT TTTGTTTTCT TTCTTTCAGC CTGAATTCTA AGCAACAAC	1018
TCAGGTTGGG GGCCTAAACT TGTTCCCTGCC TCCCTCACCC CACCCCGCCC CACCCCCAGC	1078
CCTGGCCCTT GGCTTCTCTC ACCCTCCCA AATTAAATGG ACTCCAGATG AAAATGCCAA	1138
ATTGTCATAG TGACACCAGT GGTTGTCAG CTCCTGTGCA TTCTCCTCTA AGAACTCACC	1198
TCCGTTAGCG CACTGTGTCA GCGGGCTATG GACAAGGAAG AATAGTGGCA GATGCAGCCA	1258
GCGCTGGCTA GGGCTGGGAG GGTTTTGCTC TCCTATGCAA TATTTATGCC TTCTCATTCA	1318
GAAGTGTAAAG ATGATCGCGC AGGGCATCAT GTCACCATGT CAGGTCCGGA GGGGAGGGCC	1378
TATCCCCCTA TCCCAGGCAT CCCAGACGAG GACTGGCTGA GGCTAGGCGC TCTCATGATC	1439
CACCTGCCCC GGGAGGGCAG CGGGGAAGAC AGAGAAAAGC AAAACGCATT CCTCCTCAGC	1498
TCCACCCACC TGGAGACGAA TGTAGCCAGA GAGGAGGAAG GAGGGAAACT GAAGACACCG	1558
TGGCCCCCTCG GCCTTCTCTC TGCTAGAGTT GCCGCTCAGA GGCTTCAGCC TGAATTCCAG	1618
CGGTCCCAAG AACACCTACT AATTCTTCTC CACTCCTTCA TGGCTGGGAC AGTTACTGGT	1678
TCATATGCAA GTAAAGATGA CAATTTACTC AACAAAAAAA AAAGGAATTC	1728

## (2) INFORMATION FOR SEQ ID NO:2:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 228 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met	Leu	His	Val	Glu	Met	Leu	Thr	Leu	Val	Phe	Leu	Val	Leu	Trp	Met
-19					-15			-10						-5	

- 40 -

Cys Val Phe Ser Gln Asp Pro Gly Ser Lys Ala Val Ala Asp Arg Tyr  
                     1                                    5                                    10  
 Ala Val Tyr Trp Asn Ser Ser Asn Pro Arg Phe Gln Arg Gly Asp Tyr  
                     15                                    20                                    25  
 His Ile Asp Val Cys Ile Asn Asp Tyr Leu Asp Val Phe Cys Pro His  
                     30                                    35                                    40                                    45  
 Tyr Glu Asp Ser Val Pro Glu Asp Lys Thr Glu Arg Tyr Val Leu Tyr  
                                     50                                    55                                    60  
 Met Val Asn Phe Asp Gly Tyr Ser Ala Cys Asp His Thr Ser Lys Gly  
                                     65                                    70                                    75  
 Phe Lys Arg Trp Glu Cys Asn Arg Pro His Ser Pro Asn Gly Pro Leu  
                                     80                                    85                                    90  
 Lys Phe Ser Glu Lys Phe Gln Leu Phe Thr Pro Phe Ser Leu Gly Phe  
                                     95                                    100                                    105  
 Glu Phe Arg Pro Gly Arg Glu Tyr Phe Tyr Ile Ser Ser Ala Ile Pro  
                                     110                                    115                                    120                                    125  
 Asp Asn Gly Arg Arg Ser Cys Leu Lys Leu Lys Val Phe Val Arg Pro  
                                     130                                    135                                    140  
 Thr Asn Ser Cys Met Lys Thr Ile Gly Val His Asp Arg Val Phe Asp  
                                     145                                    150                                    155  
 Val Asn Asp Lys Val Glu Asn Ser Leu Glu Pro Ala Asp Asp Thr Val  
                                     160                                    165                                    170  
 His Glu Ser Ala Glu Pro Ser Arg Gly Glu Asn Ala Ala Gln Thr Pro  
                                     175                                    180                                    185  
 Arg Ile Pro Ser Arg Leu Leu Ala Ile Leu Leu Phe Leu Leu Ala Met  
                                     190                                    195                                    200                                    205  
 Leu Leu Thr Leu

## (2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 29 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

GGATCTAGAG CACCAGCAAC ATGGATTGT

- 41 -

## (2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TCGGTCTAGA TCATTATTGG CTACTTTCAC CAGAGAT

37

## (2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 10 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Val	Asn	Phe	Asp	Gly	Tyr	Ser	Ala	Arg	Asp
1				5				10	

## (2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 16 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Val	Phe	Asp	Val	Asn	Phe	Lys	Val	Glu	Xaa	Ser	Leu	Glu	Pro	Ala	Asp
1				5				10					15		

## (2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 31 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

- 42 -

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Ala	Val	Ala	Asp	Arg	Tyr	Ala	Val	Tyr	Trp	Asn	Ser	Ser	Asn	Pro	Arg
1				5				10					15		
Phe	Gln	Arg	Gly	Asp	Tyr	His	Ile	Ile	Val	Xaa	Ile	Asn	Xaa	Tyr	
			20				25					30			

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

GAYMGNTAYG CNGTNTAYTG G

21

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 21 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

RTANCCRTCR AARTTNACCA T

21

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

- 43 -

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GGAGGACATA TGAGCCAGGA CCCGGGCTCC AAG

33

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 36 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GAAGAAGGAT CCCTATGGCT CGGCTGACTC ATGTAC

36

- 44 -

## WHAT IS CLAIMED IS:

1. A purified and isolated polypeptide capable of binding at least one EPH-like receptor  
5 wherein the receptor is the HEK4 receptor.
2. The polypeptide of Claim 1 wherein the binding results in activation of the EPH-like receptor.  
10
3. The polypeptide of Claim 1 which is human.
4. The polypeptide of Claim 1 which is  
15 substantially free of other human proteins.
5. The polypeptide of Claim 1 having in glycosylated form an apparent molecular weight of about 21-27 kD on nonreducing SDS-PAGE.  
20
6. The polypeptide of Claim 1 which is characterized by being a product of procaryotic or eucaryotic expression of an exogenous DNA sequence.
- 25 7. The polypeptide of Claim 6 wherein the exogenous DNA is cDNA, genomic DNA or synthetic DNA.
8. The polypeptide of Claim 1 which has at least about 70% homology to the amino acid sequence as  
30 shown in Figure 3 (SEQ ID NO: 1).
9. The polypeptide of Claim 1 which has the amino acid sequence of Figure 3 (SEQ ID NO: 1).
- 35 10. The polypeptide of Claim 9 further comprising a methionine residue at position -1.

11. The polypeptide of Claim 1 which is soluble or bound to a cell surface.

5 12. The polypeptide of Claim 1 which is further capable of binding the ECK receptor.

13. The polypeptide of Claim 1 which is multimeric.

10

14. The polypeptide of Claim 13 selected from the group consisting of: an immunoglobulin chimera, an antibody clustered protein, and covalently and noncovalently attached protein monomers.

15

15. An isolated nucleic acid encoding a polypeptide capable of binding to at least one EPH-like receptor wherein the nucleic acid is selected from the group consisting of:

20

a) the nucleic acid as shown in Figure 3 (SEQ ID NO: 1);

b) nucleic acids which hybridize under conditions of 6XSSC and 65°C with the coding region of Figure 3 (SEQ ID NO: 1); and

25

c) nucleic acids which are degenerate to the nucleic acids of (a) and (b).

30

16. The nucleic acid of Claim 15 which is cDNA, genomic DNA or synthetic DNA.

17. A polypeptide encoded by the nucleic acids of Claim 15.

18. The nucleic acids of Claim 15 which include one or more codons preferred for expression in Escherichia coli.

35



- 46 -

19. A biologically functional plasmid or viral vector including the nucleic acid of Claim 15.

5           20. A procaryotic or eucaryotic host cell stably transformed or transfected with a DNA vector of Claim 19.

10           21. The host cell of Claim 20 which is a CHO cell.

          22. The host cell of Claim 20 which is Escherichia coli.

15           23. A process for the production of a polypeptide capable of binding at least one EPH-like receptor comprising growing under suitable nutrient conditions procaryotic or eucaryotic host cells transformed or transfected with the nucleic acid of  
20 Claim 15 and isolating polypeptide products of the expression of the nucleic acids in the vector.

          24. A pharmaceutical composition comprising a therapeutically effective amount of the polypeptide  
25 of Claim 1 and a pharmaceutically acceptable adjuvant.

          25. The composition of Claim 24 wherein the polypeptide is of human origin.

30           26. An antibody or fragment thereof specifically binding the polypeptide of Claim 1.

          27. The antibody of Claim 26 which is a monoclonal antibody.

35

- 47 -

28. A method of activating at least one EPH-like receptor comprising contacting the receptor with the polypeptide of Claim 1.

5           29. A method of modulating the growth or differentiation of cells comprising contacting the cells with the polypeptide of Claim 1.

10           30. The method of Claim 29 wherein the cells are kidney, lung, liver, skin, digestive tract, glial, nerve, or hematopoietic cells.

15           31. A method of maintaining or restoring cellular function in the nervous system of a mammal comprising administering a therapeutically effective amount of the polypeptide of Claim 1.

20           32. A method of regenerating damaged or depleted tissue in a mammal comprising administering a therapeutically effective amount of the polypeptide of Claim 1.

25           33. The method of Claim 32 wherein the tissue is from liver, lung, digestive tract or nervous system.

30           34. A method for detecting the presence of a polypeptide capable of binding at least one EPH-like receptor in a biological sample comprising: contacting the sample with the antibody of Claim 26 under conditions allowing for antibody binding to the polypeptide; and detecting the binding of the antibody.

35           35. A method for detecting in a biological sample the presence of a nucleic acid encoding a

- 48 -

polypeptide capable of binding at least one EPH-like  
receptor comprising: contacting the sample with the  
nucleic acid of Claim 15 under conditions allowing for  
hybridization; and detecting the hybridization of the  
5 nucleic acid.

36. An anti-sense nucleotide sequence  
which hybridizes to the nucleic acid of Claim 15 in a  
manner that blocks transcription or translation of the  
10 nucleic acid.

*FIGURE 1*

BLAcore screening of conditioned media on HEK4-X surface






FIGURE 2



3 / 10

FIGURE 3

10 30 50  
GAATCCCCCAGCTTGGTGGGCGCCTCTTTCTCTCGCCCCCTTTTCATTTTTATTAT

70 90 110  
TCATATTTATTTGGCGCCCGCTCTCTCTGTCCCTTTGCCTGCCTCCCTCCCTCCGGAT

130 150 170  
CCCCGCTCTCTCCCGGAGTGGCGCGTCCGGGGCTCCGCGCTGGCCAGGCGTGATGTTG

M L  
-19

190 210 230  
CACGTGGAGATGTTGACGCTGGTGTTCCTGGTGCTCTGGATGTGTGTGTTTCAGCCAGGAC

H V E M L T L V F L V L W M C V P S Q D  
-10 -1 1

250 270 290  
CCGGGCTCCAAGGCGCTCGCCGACCGCTACGCTGTCTACTGGAACAGCAGCAACCCCGAGA

P G S K A V A D R Y A V Y W H S S N P R  
10 20

310 330 350  
TTCCAGAGGGGTGACTACCATATTGATGTCTGTATCAATGACTACCTGGATGTTTTCTGC

P Q R G D Y H I D V C I N D Y L D V F C  
30 40

370 390 410  
CCTCACTATGAGGACTCCGTCCCGAAGATAAGACTGAGCGCTATGTCTCTACATGGTG

P H Y E D S V P E D K T E R Y V L Y M V  
50 60

430 450 470  
AACTTTGATGGCTACAGTGCCTCCGACCACTTCCAAAGGGTTCAAGAGATGGGAATGT

N F D G Y S A C D H T S K G F K R W E C  
70 80

490 510 530  
AACCGGCCTCACTCTCCAAATGGACCGCTGAAGTTCTCTGAAAAATTCAGCTCTTCACT

N R P H S P N G P L K F S E K F Q L F T  
90 100

550 570 590  
CCCTTTCTCTAGGATTTGAATTCAGGCCAGGCCGAGAATATTTCTACATCTCTCTGCA

P F S L G F E F R P G R E Y F Y I S S A  
110 120

610 630 650  
ATCCAGATAATGGAAGAAGTCTCTCTAAAGCTCAAAGTCTTTGTGAGACCAACAAT

I P D N G R R S C L K L K V F V R P T N  
130 140

4 / 10

## FIGURE 3

670 690 710  
AGCTGTATGAAACTATAGGTGTTTCATGATCGTGTTCGATGTTAACGACAAAGTAGAA  
S C M K T I G V H D R V F D V N D K V E  
150 160

730 750 770  
AATTCATTAGAACCAGCAGATGACACCGTACATGAGTCAGCCGAGCCATCCCGCGCGAG  
N S L E P A D D T V H E S A E P S R G E  
170 180

790 810 830  
AACGCGGCACAAACACCAAGGATACCCAGCCGCTTTTGGCAATCCTACTGTTCTCTCTG  
N A A Q T P R I P S R L L A I L L P L L  
190 200

850 870 890  
GCGATGCTTTTGACATTATAGCACAGTCTCCTCCCATCACTTGTACAGAAAACATCAGG  
A M L L T L  
209

910 930 950  
GTCTTGGAACACCAGAGATCCACCTAACTGCTCATCCTAAGAAGGGACTTGTTATTGGGT  
970 990 1010  
TTTGGCAGATGTCAGATTTTTGTTTTCTTTCTTCAGCCTGAATTCCTAAGCAACAATTC  
1030 1050 1070  
AGGTTGGGGGCTAAACTTGTTCTGCTCCTCCCTCACCCACCCCGCCCCAGCCC  
1090 1110 1130  
TGGCCCTTGCGTCTCTCACCCCTCCCAAATTAAATGGACTCCAGATGAAAATGCCAAAT  
1150 1170 1190  
TGTCATAGTGACACCAGTGGTTCGTCAGCTCCTGTGCATTCTCCTCTAAGAACTCACCTC  
1210 1230 1250  
CGTTAGCGCACTGTGTCAGCGGGCTATGGACAAGGAAGATAGTGGCAGATGCAGCCAGC  
1270 1290 1310  
GCTGGCTAGGGCTGGGAGGGTTTTGCTCTCCTATGCAATATTTATGCCTTCTCATTGAGA  
1330 1350 1370  
ACTGTAAGATGATCGCGCAGGGCATCATGTCAACATGTGAGGTCCGGAGGGGAGGGCTTA  
1390 1410 1430  
TCCCCCTATCCCAGGCATCCCAGACGAGGACTGGCTGAGGCTAGGCGCTCTCATGATCCA  
1450 1470 1490  
CCTGCCCCGGGAGGGCAGCGGGGAAGACAGAGAAAAGCAAAACGCATTCTCTCCTCAGCTC  
1510 1530 1550  
CACCCACCTGGAGACGAATGTAGCCAGAGAGGAGGAAGGAGGGAACTGAAGACACCGTG

5 / 10


## FIGURE 3

1570 1590 1610  
GCCCCCTCGGCCTTCTCTGCTAGAGTTGCCGCTCAGAGGCTTCAGCCTGACTTCCAGCG  
1630 1650 1670  
GTCCCAAGAACACCTACTAATTCTTCTCCACTCCTTCATGGCTGGGACAGTTACTGGTTC  
1690 1710  
ATATGCAAGTAAAGATGACAATTTACTCAACAAAAAAAAAAGGAATTC



6/10

FIGURE 4



b)


*new standards*

Time Range (min)
114-117
117-120
120-123
123-126
126-129
129-132
132-135
135-138
138-141




7 / 10

FIGURE 5




8 / 10

FIGURE 6




9/10

FIGURE 7



10/10

FIGURE 8



## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07K14/47 C12N15/12 C07K16/18 C12N15/70 G01N33/68  
 C12Q1/68 A61K38/17 C07K19/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N C12Q G01N A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO,A,96 13518 (GENENTECH INC ;CARAS INGRID W (US); WINSLOW JOHN W (US)) 9 May 1996 see the whole document ---	1-35
P,X	NEURON, vol. 14, 5 May 1995, pages 973-981, XP002004532 J.W.WINSLOW E.A.: "Cloning of AL-1, aligand for an Eph-related TKR involved in axon bundle formation" see the whole document ---	1-12, 15-36
P,X	CELL, vol. 82, 11 August 1995, NA US, pages 359-370, XP002004533 U.DRESCHER E.A.: "In vitro guidance of retinal ganglion cell axons by RAGS..." see the whole document ---	1-8,11, 12,15-36
-/-		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*&\* document member of the same patent family

Date of the actual completion of the international search

3 June 1996

Date of mailing of the international search report

12.06.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+ 31-70) 340-3016

Authorized officer

Groenendijk, M

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP,A,0 597 503 (AMGEN INC) 18 May 1994 cited in the application see the whole document ---	1-36
A	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 89, March 1992, WASHINGTON US, pages 1611-1615, XP002004534 I.P.WICKS E.A.: "Molecular cloning of HEK..." cited in the application see the whole document -----	1-36

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US96/01079

**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claims 28-33 encompass or are directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the compound/composition. ./...
2. ☐ Claims Nos.:  
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

**Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)**

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.



FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

Annex:

The peptides of the claims 1-7 are essentially only characterised by being "capable of binding at least one EPH-like receptor wherein the receptor is the HEK4 receptor". In the absence of any structural feature these peptides are considered to be ill-defined and the claims 1-7, 11-14 and the related claims 24-34 are considered to contravene Art.6 PCT, not allowing a complete search (Art.17(2)(a)(ii)PCT).

As far as the peptides of the present application are concerned the search has been directed to the peptides defined in claim 8 and fragments thereof.

Incomplete search:

Claims searched completely: 9-10, 15-23, 35, 36.

Claims searched incompletely: 1-7, 11-14, 24-34.

## INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/01079

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9613518	09-05-96	NONE	
EP-A-0597503	18-05-94	AU-B- 5599794	08-06-94
		CN-A- 1094446	02-11-94
		FI-A- 952328	10-07-95
		NO-A- 951861	11-07-95
		WO-A- 9411020	26-05-94